



Age and Time-Dependent Risk Model Associated with Progressing to Dementia

Kaur, D., Bucholc, M., Finn, D., Todd, S., Wong-Lin, K., & McClean, P. (2019). Age and Time-Dependent Risk Model Associated with Progressing to Dementia. In *Alzheimer's and Dementia* (7 ed., Vol. 15, pp. 237-238). Elsevier. <https://doi.org/10.1016/j.jalz.2019.06.043>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Alzheimer's and Dementia

Publication Status:
Published (in print/issue): 14/07/2019

DOI:
[10.1016/j.jalz.2019.06.043](https://doi.org/10.1016/j.jalz.2019.06.043)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Age and time-dependent risk model for progressing to dementia

Kaur, D.¹, Bucholc, M.², Finn, D.P.³, Todd, S.⁴, Wong-Lin, K.², McClean, P.L.¹

¹ Northern Ireland Centre for Stratified Medicine, Biomedical Sciences Research Institute, Derry~Londonderry, Northern Ireland, UK

² Intelligent Systems Research Centre, School of Computing, Engineering and Intelligent Systems, Ulster University, Derry~Londonderry, Northern Ireland, UK

³ Pharmacology and Therapeutics, School of Medicine, National University of Ireland Galway, Republic of Ireland

⁴ Altnagelvin Area Hospital, Western Health and Social Care Trust, Derry~Londonderry, Northern Ireland, UK

Background: This study evaluated some of the major cardio-metabolic risk factors associated with progression to dementia, among different age groups in healthy and MCI subjects. These risk factors were also analysed with respect to duration of progression.

Methods: The NACC database, one of the largest, most comprehensive longitudinal databases for dementia research was used in this study. Risk factors analysed, using multivariate-logistic regression were age, gender, smoking, BMI, alcohol-abuse, stroke, heart-attack/cardiac-arrest, geriatric depression scale (GDS), medications, diabetes, hypertension and hypercholesterolemia. Analyses were conducted at baseline and over time in those who progressed from healthy-to-dementia, and from MCI-to-dementia. Combinations of variables were also evaluated and ranked on basis of increasing risk. Analyses were stratified by age, gender and duration of progression. The aim is to generate a risk model, applicable across the life-course, to predict risk of progression from healthy and MCI to dementia.

Results: Active diabetes increased the risk of progression from healthy to dementia (odds ratio- OR: 3.274, p: $6e^{-04}$) in 61-70 age group. Individuals aged 81-90 years with active hypercholesterolemia were at reduced risk of progressing from healthy to dementia (OR: 0.59, p: 0.0023). Active hypertension reduced the risk of progression from MCI to dementia (OR: 0.8091, p: 0.0051) in 71-80 age group.

Depression (GDS score 6-10) in general, was significantly associated with increased risk of progression from healthy to dementia, irrespective of the duration of progression. Medications reduced the risk of progressing to dementia within 4 years, in the 71-80 age group (OR: 0.139, p: 0.0026). Healthy individuals aged 81-90 with active diabetes were at increased risk of progressing to dementia within 4 years (OR: 5.1798, p: 0.0012). Additionally, healthy individuals aged 61-70 with active diabetes, were at increased risk of progression (OR: 22.267, p< $2e^{-04}$), between 8~12 years from baseline.

High BMI in general (≥ 30) was significantly associated with reduced risk of MCI to dementia progression, over ~12 years from baseline. MCI patients with active hypertension were at reduced risk of progressing (OR: 0.8114, p: 0.0022) within 4 years from baseline. MCI patients aged 81-90 with a history of stroke, were at increased risk of progressing to dementia in ≤ 4 years (OR: 1.9052, p: 0.0076). Those aged ≥ 91 with GDS score 6-10 were also at increased risk of progressing to dementia within 4 years (OR: 9.8376, p: $4e^{-04}$).

Conclusions: Different risk factors are particularly relevant at different ages, and in terms of duration of progression to dementia, from healthy and MCI stages.